Information Sheet


IS13091

Canegrub identification: Central Region

Anal hair (raster) patterns of canegrubs found in the Plane Creek to Proserpine region:







1. Greyback


Two almost straight rows of 20-28 short hairs.


2. Squamulata

Two straight rows of 28-40 short thick hairs, wider gap at front.

3. Grata

Two slightly curved rows of 18-26 thick hairs, hairs in each row close together compared with greyback.





Further information

For help with identifying canegrub species or further information please contact your local adviser or SRA on 3331 3333, or visit www.sugarresearch.com.au

4. French's

Pear-shaped, about 50 long hairs each side.

5. Negatoria

Pear-shaped, about 50 long hairs each side.

sugarresearch.com.au

Copyright © 2013 * All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior permission of SRA. Disclaimer in this disclaimer a reference to 'we', 'us' or 'or 'or' means SRA and our directors, officers, agents and employees. Although we do our best to present information that is correct and accurate, we make no warranties, guarantees or representations about the suitability, reliability, reliability, reliability, reliability, reliability, and accurate, or accurately of the information of the persent in this information appearing in this Information sheet, and you agree that we will not be liable for any loss or damage whatsoever (including through negligence) arising out of, or in connection with the use of this Information sheet. We recommend that you contact our staff before acting on any information provided in this Information Sheet. Warning Our tests, inspections and recommendations should not be reliad on without current, when you not be accurate, complete or applicable for your particular needs for many reasons, including (for example) SRA being unaware of other matters relevant to individual crops, the analysis of unrepresentative samples or the influence of environmental, managerial or other factors on production.